

# Arctic sea ice decline weakens the Atlantic meridional overturning circulation

Florian Sévellec


Ocean and Earth Science, University of Southampton

A. V. Fedorov & W. Liu - Yale University

Observing and Understanding the AMOC  
May 2017

## Is the AMOC slowing down?


◇ Rapid measurement of AMOC at 26°N:



Smeed et al. (2014)

⇒ **The AMOC seems to slow-down on decadal timescales.**

# Is the AMOC slowing down on longer timescale?



⇒ This "Warming Hole" has been conjecture to be the signature of a **long-term AMOC slow-down.**


(Drijfhout et al., 2012; Rahmstorf et al., 2015)

NASA GISS temperature data.

◇ Objective: **Can we attribute this AMOC slow-down?**

◇ Method:


1. We compute AMOC sensitivity to constant surface buoyancy fluxes,
  - ▶ Optimization procedure based on Lagrangian multipliers,
  - ▶ This requires the “adjoint” model (NEMO-OPA at  $2^\circ$  resolution),
  - ▶ AMOC - baroclinic volume transport at 1,500 m depth and  $50^\circ$ N.
2. Optimal fluxes for a range of imposed duration from 10 yr to 200 yr.
3. We separate the fluxes in 3 regions:



- ▶ **ARCT**: Arctic Ocean,
- ▶ **NATL**: North Atlantic,
- ▶ **ROTO**: Rest Of The Oc.

⇒ **A dynamical attribution of heat flux influences on the AMOC.**

# AMOC sensitivity to constant heat flux




- ▶ **North Atlantic** dominates changes from years to a few decades,
- ▶ **Arctic** dominates changes from several decades to centuries.

(Fluxes are normalized to  $1 \text{ W m}^{-2}$ .)

⇒ On centennial timescales, despite NATL being 3 times as large as ARCT, ARCT is twice as effective for weakening the AMOC!

# AMOC sensitivity to constant heat flux

OPTIMAL HEAT FLUX PERTURBATION OF AMOC INTENSITY ( $W m^{-2}$ )



(Fluxes are normalized to  $1 W m^{-2}$  for comparison.)


- ▶ On **annual to decadal timescales**, the sensitivity is located in the **subpolar N. Atl.**,

- ▶ On **multidecadal timescales**, the sensitivity further expand northward, spreading in the **Arctic region**.

# Change in oceanic heat flux in the Arctic since 1979

## RECONSTRUCTED NET OCEANIC HEAT FLUX CHANGES

[2005-2014]  
vs  
[1979-1988]




◇ ERA-Interim structure,

◇ Uniformly rescaled to  $+0.55 \text{ W m}^{-2}$  (IPCC, 2013).

⇒ Arctic is warmed at a rate of  $\sim 8 \text{ W m}^{-2}$  over the last 35 yr.

# AMOC response to change in surface oceanic heat flux

- ▶ A 100-yr steady change, equivalent to the current one, would slow down the AMOC by 4 Sv,




- ▶ The **Arctic** contribution to this slow-down is nearly **3 times as efficient** as the **N. Atl.** one.

- ▶ Unlike for other oceanic regions, **Arctic** heat flux change **reduces northward Atl. heat transport**  $\Rightarrow$  *Warming Hole?*





## Change in Arctic sea ice cover since 1979

ANNUAL SEA ICE CONCENTRATION (%)




- ▶ There is a **reduction** of the **sea ice cover** in the Arctic since 1979,



- ▶ This **increases** the **heat flux** reaching the ocean surface.


# Simulating Arctic sea ice changes




## A coupled experiment (CESM):

- ▶ **Modifying** the properties of sea ice and snow:
  - ▶ Reduce the emissivity (LW),
  - ▶ Reduce the albedo (SW).
- ▶ Reproducing quantitatively the observed **Arctic sea ice decline**.

# Impact of Arctic sea ice change in the North Atlantic



**(d) SST ANOMALIES (°C)**  
**LW**



- ▶ **AMOC slow-down** by  $\sim 50\%$  in 100 yr,
- ▶ **Cooling of up to  $2^\circ\text{C}$**  in the subpolar N. Atl.

## Conclusion

### Attribution of the AMOC slow-down to buoyancy fluxes.

- ▶ Potential impacts:
  - ▶ **Subpolar North Atlantic** has higher potential impact on timescales from **years to a few decades**,
  - ▶ **Arctic** has higher potential impact on timescales from **several decades to centuries** (6 times as efficient).
  - ▶ Results for freshwater are qualitatively equivalent but quantitatively one order of magnitude weaker than for heat.

### Reduction in Arctic sea ice increases ocean heat uptake.

- ▶ Projected impacts:
  - ▶ **AMOC** would **slow-down by 4-6 Sv in a century** under current observed changes,
  - ▶ **30%** can be attributed to the **N. Atl.** and **60%** – to the **Arctic**.
  - ▶ Consequently, N. Atl. **STT** would **cool-down by up to 2°C**.